# **Dr. Gustav Fischer Before Surgery Instructions**

### MULTNOMAH ORTHOPEDIC CLINIC 503-231-1426

#### **Before Surgery**

- 1. STOP aspirin 7 days before surgery
- 2. STOP blood thinners:

Coumadin - 5 days before surgery

Plavix - 5 days before surgery

Eliquis, Pradaxa, Xarelto - 3 days before surgery

- 3. STOP SGLT2s (Jardiance, Farxiga, Invokana) 3 days before surgery.
- 4. STOP GLP-1RAs (Ozempic, Wegovy, Mounjaro, etc.) 1 week before surgery
- 5. STOP NSAIDs (e.g. advil, aleve, celebrex, mobic) 7 days before surgery
- 6. STOP shaving or waxing legs 7 days before surgery
- 7. STOP ACE inhibitors and ARBs the day before surgery
- 8. STOP herbal supplements 7 days before surgery
- 9. Continue Beta blockers the morning of surgery with a small sip of water
- 10. Drink plenty of water the day before surgery
- 11. Nothing to eat or drink after midnight
- 12. If you do not have a walker, arrange for one prior to surgery

#### **Day of Surgery**

- 1. Bring photo ID, insurance card and glasses if needed for filling out paperwork
- 2. NO jewelry, eye make-up, or contact lenses
- 3. Please wear loose, comfortable clothing
- 4. Plan to have transportation and 24 hour care after discharge
- 5. Bring your walker with you to surgery

### **After Surgery Medications**

## Prescribed medications below should be filled prior to surgery and only taken after surgery Naproxen 250mg twice daily for 14 days Naproxen 500mg twice daily for 14 days

- Tylenol 1000 mg every 6 hours for 14 days
- ' Aspirin 81 mg twice daily for 42 days
- Eliquis 2.5 mg twice daily for 7 days, then resume 5 mg twice daily
- Tranexamic Acid (TXA) 1950 mg daily with breakfast for 3 days
- Zofran 4 mg 1 tablet every 6 <u>as needed</u> for nausea
- Tramadol 50mg 1-2 tablets every 6 hours <u>as needed</u> for pain
- Oxycodone 5mg 1-2 tablets every 4 hours <u>as needed</u> for pain
- Oxycodone 5mg 1-3 tablets every 3 hours <u>as needed</u> for pain
- Dilaudid 2mg 1-3 tablets every 3 hours <u>as needed</u> for pain
- Cefdinir 300 mg twice daily for 7 days
   ☐ Doxycycline 100mg twice daily for 7 days